

Name (Print) _____ Date _____

Grand Rounds Enduring Materials
Diagnosis and Management of the Pediatric Neck Mass

Please select the best answer:

1. What percentage of neck masses in children is benign?
 - a. 20%
 - b. 40%
 - c. 70%
 - d. 90%

2. What percentage of pediatric neck masses, that undergo biopsy, are congenital in etiology?
 - a. 35%
 - b. 55%
 - c. 75%
 - d. 95%

3. A malignant pediatric neck mass in the age range of 7 – 13 years, is most likely due to what?
 - a. Neuroblastoma
 - b. Rhabdomyosarcoma
 - c. Hodgkin's Lymphoma
 - d. Thyroid cancer

4. A pediatric neck mass associated with enlargement and pain while eating is most likely associated with what etiology?
 - a. Sialadenitis
 - b. Venous malformation
 - c. Hemangioma
 - d. Hematoma

5. What location is typically associated with a branchial cleft cyst?
 - a. Anterior deep cervical
 - b. Submandibular triangle
 - c. Midline
 - d. Submental

In order to receive CME credit, please complete the program evaluation and the questions.

These can be emailed to Linda Carpenter at:

LCarpenter@svhs.org

or faxed to 814-452-7131

or mailed to:

Linda Carpenter
CME Coordinator
Saint Vincent Health Center
232 West 25 Street
Erie, PA 16544

If you have any questions, please contact Linda Carpenter at 814-452-5717.

Thank you

SAINT VINCENT HEALTH CENTER
CME DEPARTMENT